Streamflow and Groundwater Conditions in Rhode Island

October 22, 2020

C. Bruet
U.S. Geological Survey
New England Water Science Center



Table 2, Rhode Island Drought Indices and Phases

Drought Phase	Palmer Drought Index +	Crop Moisture Index	Precipitation +	Ground Water** +	Stream flow +	Reservoirs**
Normal	-1.0 to -1.99	0.0 to -1.0	Slightly Dry	1 month below normal 1 month below normal	2 consecutive months below normal	Reservoir levels at or near normal for the time of year
Advisory	-2.0 to -2.99	-1.0 to -1.9 Abnormally Dry	2 month cumulative below 65% of normal	At least 2 out of 3 months below normal	3 consecutive months below normal	Small index Reservoirs below normal
Watch	-3.0 to -3.99	-2.0 to -2.9 Excessively Dry	1 of the following criteria met: 3 month cum. <65% or 6 month cum. <70% or 12 month cum. <70%	4-5 consecutive months below normal	At least 4 out of 5 consecutive months below normal	Medium index Reservoirs below normal
Warning	-4.0 and below	> -2.9 Severely Dry	2 out of 3 of the above criteria met: 3 month cum. <65% and 6 month cum. <65% or 6 month cum. <65% and 12 month cum. <65% or 3 month cum. <65% and 12 month cum. <65% and	6-7 consecutive months below normal observation wells recording monthly record lows	At least 6 out of 7 consecutive months below normal	Large index reservoirs below normal
Emergency	-4.0 and below	> -2.9 Severely dry	Same criteria as Warning and Previous month was Warning or Emergency	>7 months below normal Observation wells recording monthly record lows	>7 months below normal	Continuation of previous month's conditions

⁺ Major Hydrologic Indicators.

[&]quot;Normal" is defined as the statistical average of the data for the period of record. Percentages for precipitation are relative to normal.




^{**} Local triggers from the water system supply management plans will also be considered in a assessing drought phases on a regional basis. The WRB staff will review local plans and work with suppliers to coordinate regarding drought phases and to collect, review and report surface reservoir and ground water data.

Average Streamflow Conditions July – September











Percent by Drought Region

Drought Region	Number of	June Avg	July Avg	August Avg	September Avg	October 1-19 Avg
Drought Region	Gages	Percentile	Percentile	Percentile	Percentile	Percentile
Central East	4	24.32	25.25	12.52	4.58	10.25
Central West	1	15.15	36.36	3.03	3.03	3.23
North East	4	24.50	35.03	10.26	10.27	16.86
North West	1	44.64	62.50	7.14	5.36	3.57
Southern	11	28.57	26.22	8.53	4.94	3.90

N-Day Data as of October 20, 2020

Drought Region	Number of Gages	7-Day Avg Percentile	14-Day Avg Percentile	28-Day Avg Percentile
Central East	4	53.65	36.24	15.62
Central West	1	32.36	16.09	10.52
North East	4	57.54	40.72	23.32
North West	1	16.35	7.70	3.19
Southern	11	29.45	13.98	6.42




Statewide Average Percentages


June Avg	July Avg	August Avg	September Avg	October 1-19
Percentile	Percentile	Percentile	Percentile	Avg Percentile
27.11	29.92	9.29	5.82	7.53
Low End of	Low End of	Much Below	Much Below	Much Below
Normal	Normal	Normal	Normal	Normal

Explanation - Percentile classes							
Low	<10	10-24	25-75	76-90	>90	Lliab	
Low	Much below normal	Below normal	Normal	Above normal	Much above normal	High	




N-Day Maps as of 10/20/2020






Current Streamflow Conditions – October 21, 2020

Explanation - Percentile classes							
Low	<10	10-24	25-75	76-90	>90	Lliab	
LOW	Much below normal	Below normal	Normal	Above normal	Much above normal	High	










science for a changing world


Last updated: 2020-10-21






science for a changing world

Last updated: 2020-10-21








science for a changing world


Last updated: 2020-10-21

Groundwater Conditions



science for a changing world

415546071474701 - RI-BUW 395 BURRILLVILLE, RI



Plot created 10/20/20 21:17


Explanation - Percentile Classes

10 - 24 25 - 75 76 - 90


Monthly Median




Data Point











Groundwater Conditions

Well	Tape down
LTW 142	21.33
POW 551	10.9
COW 342	20.02
WCW 59	DRY
NSW 21	11.96
CUW 265	15.88






<10 10-24 25-75 76-90 >90


Monthly Median




Data Point












Note: The most recent measurement on 10/16/2020 has the following status: 'The site was dry (no water level is recorded).'






Table 2, Rhode Island Drought Indices and Phases

Drought Phase	Palmer Drought Index +	Crop Moisture Index	Precipitation +	Ground Water** +	Stream flow +	Reservoirs**
Normal	-1.0 to -1.99	0.0 to -1.0	Slightly Dry	1 month below normal 1 month below normal	2 consecutive months below normal	Reservoir levels at or near normal for the time of year
Advisory	-2.0 to -2.99	-1.0 to -1.9 Abnormally Dry	2 month cumulative below 65% of normal	At least 2 out of 3 months below normal	3 consecutive months below normal	Small index Reservoirs below normal
Watch	-3.0 to -3.99	-2.0 to -2.9 Excessively Dry	1 of the following criteria met: 3 month cum. <65% or 6 month cum. <70% or 12 month cum. <70%	4-5 consecutive months below normal	At least 4 out of 5 consecutive months below normal	Medium index Reservoirs below normal
Warning	-4.0 and below	> -2.9 Severely Dry	2 out of 3 of the above criteria met: 3 month cum. <65% and 6 month cum. <65% or 6 month cum. <65% and 12 month cum. <65% or 3 month cum. <65% and 12 month cum. <65% and	6-7 consecutive months below normal observation wells recording monthly record lows	At least 6 out of 7 consecutive months below normal	Large index reservoirs below normal
Emergency	-4.0 and below	> -2.9 Severely dry	Same criteria as Warning and Previous month was Warning or Emergency	>7 months below normal Observation wells recording monthly record lows	>7 months below normal	Continuation of previous month's conditions

⁺ Major Hydrologic Indicators.

[&]quot;Normal" is defined as the statistical average of the data for the period of record. Percentages for precipitation are relative to normal.



^{**} Local triggers from the water system supply management plans will also be considered in a assessing drought phases on a regional basis. The WRB staff will review local plans and work with suppliers to coordinate regarding drought phases and to collect, review and report surface reservoir and ground water data.

Summary: Streamflow -

- Based on the State Guide Plan 721
 - Drought advisory calls for three consecutive months of below normal streamflow
 - August to October (through 10/20)
 - August was much below normal statewide (9.29%).
 - September was much below normal statewide (5.82%).
 - October 1 -19 has been much-below normal statewide (7.53%).
 - Oct 13-20 (7-day) has shown improvement with the two rain events on Oct 13 and Oct 16/17 (38.92%).
 - Streamflow recovery/decline will be dependent on the precipitation going forward through October and beyond.



Summary: Ground Water -

Based on the State Guide Plan 721

 Drought advisory calls for two out of three months of below normal water levels

Statewide the trigger –

- July to September <u>does meet</u> the trigger for a drought advisory
 - July central west and eastern regions were below normal. Rest of state was normal.
 - August and September were both much-below normal.
- August to October (though Oct 20)
 - The conditions remain much-below normal statewide.
- Groundwater recovery/decline will be dependent on the precipitation going forward through October and beyond.



Questions?

